

Ovine mitochondrial DNA: mapping and sequencing data in comparison with bovine mtDNA

S Hiendleder, W Hecht, R Wassmuth

*Institut fuer Tierzucht und Haustiergenetik
der Justus-Liebig-Universitaet, Gießen, Germany*

(Proceedings of the 9th European Colloquium on Cytogenetics of Domestic Animals;
Toulouse-Auzeville, 10-13 July 1990)

mtDNA / evolutionary relationship / sheep / cow

INTRODUCTION

Mitochondrial DNA (mtDNA) is maternally inherited (Gyllensten *et al*, 1985) and evolves 5-10 times faster than single-copy nuclear DNA in mammals (Brown *et al*, 1979). These properties make mtDNA an excellent tool for analyzing evolutionary relationships (Wilson *et al*, 1985).

We studied the mtDNA structure of *Ovis aries* and compared it to the *Bos taurus* (Anderson *et al*, 1982) sequence.

MATERIALS AND METHODS

mtDNA was prepared from the liver of a single Merinolandschaf, using the method described by Hecht *et al* (1988). Restriction endonucleases (*EcoRI*, *EcoRV*, *BamHI*, *BglII*, *HindIII*, *HincII*, *KpnI*, *PstI* and *XbaI*) were used according to the manufacturers' recommendations. DNA fragments were separated in 1% agarose gels, stained with ethidium bromide, photographed and transferred onto Hybond-N-Nylon membranes (Amersham product information). Hybridization analysis was performed using a [³²P]dCTP-labeled (Sambrook *et al*, 1989) 2.5 kb *HindIII* fragment of porcine mtDNA, encompassing parts of the ATPase6 and ND4 genes and the complete COIII, ND3 and ND4L genes. Two 2.6 and 3.4 kb *EcoRI* fragments of ovine mtDNA were isolated on diethylaminoethyl (DEAE) membrane filters, cloned into puc13 and the ligation products used to transform *E coli* (JM83) (Sambrook *et al*, 1989). Plasmid DNA for dideoxysequencing was prepared as described in the Diagen Application Protocol (1988). ³⁵S-Sequencing reactions were performed with the T₇-sequencing kit supplied by Pharmacia according to the manufacturers' instructions. Cronex X-ray film (Dupont) was used for autoradiographs.

RESULTS AND DISCUSSION

Total ovine mtDNA as well as cloned fragments were used to map 28 cleavage sites produced by *EcoRI*, *EcoRV*, *BamHI*, *BglII*, *HindIII*, *HincII* and *XbaI* by the double-digestion method. Hybridization analysis with a fragment of porcine mtDNA of known gene content along with sequencing data was used for orientation of the map along the bovine sequence (Anderson *et al*, 1982). The 9 enzymes (*PstI* and *KpnI* apparently do not cut ovine mtDNA) used, have a total of 31 recognition sites in bovine mtDNA (there is no *EcoRV* site in bovine mtDNA). Of the 28 restriction sites observed in the ovine sequence, only 11 are assumed to be present in the bovine sequence (fig 1). The proportion of sites shared is expected to decline as the organisms' DNA sequences diverge. The application of equations 10 and 8 from Nei and Li (1979) to these data yields an estimated sequence divergence of 16.4% between the two species.

Sequencing analysis of the terminals of 2 cloned fragments yielded 1.105 kb of sequence information from ovine mtDNA. Genes partially sequenced were ND5, Cyt.b, ATPase6 and COIII. The comparison of nucleotide and predicted amino acid sequences of the 2 species revealed a clustering of replacement nucleotide substitutions within parts of the sequenced regions of the ND5 (fig 2), Cyt.b and COIII genes (data not shown). Silent substitutions on the other hand seem to be distributed evenly across the sequences. Gene-specific and overall sequence differences are given in table I. The 4 genes showed almost the same rate of nucleotide substitutions, although the COIII (76-79%) and Cyt.b genes (73-74%) were more conserved than the ND5 gene (65-71%), for example, when compared between mouse, cow and human (Anderson *et al*, 1981, 1982; Bibb *et al*, 1981). This was probably a result of partial analysis of the genes. Amino acid replacements were highest in the ND5 gene. The lack of any amino acid replacements in the short stretch of ATPase6 DNA was probably due to the high conservation of this region within the group of species mentioned above.

Table I. Sequence differences between ovine and bovine mtDNAs at the nucleotide and predicted amino acid levels.

Gene	Nucleotides sequenced	% of gene	Nucleotide substitutions		Amino acid substitutions	
			n	%	n	%
ATPase6	74	10.9	11	14.9	0	0
COIII	193	24.7	30	15.5	5	7.8
Cyt.b	279	24.6	43	15.4	6	6.5
ND5	559	30.8	86	15.4	16	8.6
total	1105	6.7 ^a	170	15.4	27	7.3

^a) % of mt genome.

More than 80% of all nucleotide replacements were transitions. This is in accordance with results obtained by Wilson *et al* (1985) in primates.

The overall sequence difference of 15.4% at the nucleotide level corresponds well with 16.4% sequence divergence calculated from restriction map comparison. Using the data reported by Wilson *et al* (1985), sheep and cow would have evolved from a common ancestor $7.7-16.4 \times 10^6$ years ago.

An example illustrating the limitations inherent in the comparison of restriction mapping data is a *Bgl*III site, assumed to be identical in the ovine and bovine maps, which corresponds to bovine nucleotide no 12 706 according to Anderson *et al* (1982). The recognition site in the bovine sequence is actually not identical to the ovine site as shown in figure 1, but 5 nucleotides downstream from it. Taking this into account, the calculated sequence divergence is 18% according to Nei and Li (1979), which results in $9.0-18.0 \times 10^6$ years as the time of divergence between the two species.

REFERENCES

- Anderson S, Bankier AT, Barrell BG, De Bruin MHL, Coulson AR, Drouin J, Eperon IC, Nierlich DP, Roe BA, Sanger F, Schreier PH, Smith AJH, Staden R, Young IG (1981) Sequence and organization of the human mitochondrial genome. *Nature* 290, 457-464
- Anderson S, De Bruijn HL, Coulson AR, Eperon IC, Sanger F, Young IG (1982) Complete sequence of bovine mitochondrial DNA. Conserved features of the mammalian mitochondrial genome. *J Mol Biol* 156, 683-717
- Bibb MJ, Van Etten RA, Wright CC, Walberg MW, Clayton DA (1981) Sequence and gene organization of mouse mitochondrial DNA. *Cell* 26, 167-180
- Brown WM, George M, Wilson AC (1979) Rapid evolution of animal mitochondrial DNA. *Proc Natl Acad Sci USA* 76, 1967-1971
- Gyllensten U, Wharton D, Wilson AC (1985) Maternal inheritance of mitochondrial DNA during backcrossing of two species of mice. *J Hered* 76, 321-324
- Hecht W, Hiendleder S, Reis A, Dzapo V, Wassmuth R, Herzog A (1988) Analysis of mtDNAs from sheep and pig. In: *Proc of the 8th European Colloquium on Cytogenetics of Domestic Animals, Bristol, 19-22 July*
- Nei M, Li WH (1979) Mathematical model for studying genetic variation in terms of restriction endonucleases. *Proc Natl Acad Sci USA* 76, 5269-5273
- Sambrook J, Fritsch EF, Maniatis T (1989) *Molecular Cloning*. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY
- Wilson AC, Cann RL, Carr SM, George M, Gyllensten UB, Helmbychowski KM, Higuchi RG, Palumbi SR, Prager EM, Sage RD, Stoneking M (1985) Mitochondrial DNA and two perspectives on evolutionary genetics. *Biol J Linn Soc* 26, 375-400