

**Present stage of embryo transfer
and the International Embryo Transfer Society (I.E.T.S.)**

W.W. LAMPETER

*Institut für Tierzucht und Tierhygiene der L.M.U.,
8000 München 22, Federal Republic of Germany*

The *International Embryo Transfer Society* (I.E.T.S.) has 560 members from 26 countries (Jan. 1982) and is serving its members with information related to E.T. This service includes abstracts, literature reviews and summaries, published in a newsletter. It holds a yearly meeting and publishes all reports presented at the meeting in « *Theriogenology* ».

About 35 000 calves were produced worldwide by E.T. techniques in 1981. Embryo collection and transfer of bovine embryos is done worldwide mostly non-surgically. The most widely used farm animal for E.T. is the cow.

Cow embryo culture *in vitro*

L.K. ERNST, A.K. GOLUBEV, Z.N. MAKAROVA, R.S. MAMLEEV
and T.I. KUZMINA

All-Union Research Institute of Farm Animal Breeding and Genetics, Leningrad, U.S.S.R.

Fertilization of the cow oocytes matured *in vitro* to the metaphase II stage resulted in the cleavage of the oocytes. The percentage of cleaving oocytes averaged 13.9. In some experiments it amounted to 33.7 p. 100. The cleaving embryos developed to the morula and even blastocyst stage. The mechanisms of the cleavage of cow oocytes are discussed.

**Immuno- and cytogenetic methods and distant hybridization
in improvement of animals**

V.N. TIKHONOV

*Institute of Cytology and Genetics, Academy of Sciences of the U.S.S.R.,
Siberian Division, Novosibirsk, U.S.S.R.*

The new forms of commercial *LWH* (*Landrace* × *Wild Boar Hybrid*) are the result of hybridization between *Swedish Landrace* and *European, Middle Asian* and *Far East* wild boars. They have good meat features and a high resistance to unfavourable conditions of industrial pig breeding. About 3000 *LWH* of $F_1 - F_8$ were studied by immuno- and cytogenetic methods. A wide range of antigens and 5 identified chromosomes were used as genetic markers. The *LWH* homo- and heterozygous for translocations 15/17 and 16/17 ($2n = 36$ and 37) with high viability and fertility were produced.