

Isolation of subtelomeric DNA sequences labelling sheep and goat chromosome ends

Daniel VAIMAN^a, Ana BRUNIALTI^a,
Mohamed BENZAADA^b, Céline DERBOIS^a, Anne VAIMAN^a,
Allan CRAWFORD^c, Philippe METEZEAU^b, Edmond P. CRIBIU^{a,*}

^a Laboratoire de génétique biochimique et de cytogénétique,
Département de génétique animale, Institut national de la recherche agronomique,
78352 Jouy-en-Josas Cedex, France

^b Laboratoire de cytométrie analytique, Institut Pasteur, 25 rue du Dr Roux,
75014 Paris, France

^c AgResearch Molecular Biology Unit, University of Otago, P.O. Box 56 Dunedin,
New Zealand

(Received 3 May 2000; accepted 15 September 2000)

Abstract – Two techniques that make it possible to isolate telomere DNA are presented, using sheep as an example. The first technique is based upon the screening of a sheep BAC library with PCR amplified DNA segments preserved from high-power laser beam irradiation. Twenty-three BACs hybridising to 13 subtelomeric regions in sheep and goats were obtained (out of 27 in the sheep complement), of which 13 recognised more than one region, telomeric or not. Twenty-three microsatellites were isolated from these BACs and 22 were genetically mapped on the sheep international genetic map, always consistently with the cytogenetical localisation in 17 cases out of 22. These results are discussed. The second technique is based upon the selective cloning of subtelomeric enriched DNA. Preliminary results were obtained by this approach.

telomeres / FISH / sheep / goats

Résumé – Isolement de séquences d'ADN subtélomériques chez les ruminants. Deux techniques permettant d'isoler l'ADN télomérique sont présentées, le mouton servant d'exemple. La première technique est basée sur le criblage d'une banque de BACs ovins avec des segments d'ADN amplifiés par PCR préservés de l'irradiation par un laser à forte puissance. Vingt trois BACs s'hybridant à 13 régions sub-télomériques (parmi les 27 du génome ovin) ont été obtenus, dont treize reconnaissent plus qu'une région unique, télomérique ou non. Vingt-trois microsatellites ont été isolés à partir des BACs et 22 cartographiés génétiquement sur la carte ovine internationale, de façon cohérente avec la localisation cytogénétique dans 17 cas sur 22. Ces résultats sont discutés. La seconde technique est basée sur le clonage

* Correspondence and reprints
E-mail: cribiu@biotec.jouy.inra.fr

sélectif d'ADN enrichi en régions sub-téломériques. Des résultats préliminaires ont été obtenus par cette approche.

téломères / FISH / ovin / caprin

1. INTRODUCTION

Chromosome ends constitute a complex problem for pluricellular organisms. The reason for this lies in DNA replication, which, after unwinding the DNA helix is initiated with an RNA primer followed by the polymerisation of an Okazaki fragment. This RNA primer is degraded later, locally leaving a single-stranded DNA molecule. Inside a chromosome, DNA polymerase fills the hole that is generated and the extremities are ligated; in contrast to the mammalian chromosome end, a single strand overhang of 50–100 nucleotides persists [11]. Therefore, at each cell division, the telomere end shortens, at least in all somatic and non-cancerous cells, where the telomerase gene is not expressed. The mammalian chromosome end consists of 10 to 20 kb of TTAGGG repeats — very recently shown to be organised in a loop structure [9] — whose length varies according to the species and the number of cell generations since birth. Most knowledge about mammalian telomeres has been collected in mice and humans. However, it would be interesting to obtain data about telomeric regions in other mammalian species, particularly those of economic interest such as ruminants, where recently developed genomic tools and applications could benefit from such knowledge. For instance, the construction of genetic maps of ruminants has been an intense activity of animal geneticists in the last few years [2, 4, 19]. However, the true coverage of linkage groups along chromosome length has been addressed only statistically, and no tool is truly available today to measure the distance between the last marker of a linkage group and the physical chromosome end. Furthermore, the development of telomeric probes could help assess the role of cryptic chromosome translocations associated with infertility in sheep. Finally, since the evolution of mammalian genomes involves the fusion and fission of ancestor chromosome segments, telomeric probes developed in one species could be hybridised in another to detect remnants of ancient telomeric regions inside chromosomes. Since the interspecific hybridisation of large DNA probes such as BACs or YACs work well, such an approach would be easy to implement, as long as the probes are available.

Ruminant telomeric sequences are composed of the TTAGGG paradigm sequence [10], present at the end of all chromosomes. By contrast, subtelomeric sequences have been shown in humans to be characteristic of one or a few chromosomes [7, 8], suggesting that the same situation could be observed in other species. Until now, however, to the best of our knowledge, no effort has been devoted to clone this type of subtelomeric chromosomal regions in domestic species. In this study, our aim was to develop molecular techniques making it possible to isolate such sequences in sheep. Two original approaches have been elaborated and will be presented, together with some preliminary results suggesting that as in humans, subtelomeric sequences are present on a few chromosomes in ruminants. The telomeric BACs obtained were subcloned in order to isolate microsatellites. These new markers were incorporated on the international genetic map, making it possible to ascertain the genetic position of the markers.

2. MATERIALS AND METHODS

2.1. Cell lines

A primary sheep cell line SBK (54, XY) was used to prepare metaphase chromosomes prior to laser irradiation. Cells were grown to confluency in RPMI medium with 10% foetal calf serum, and plated at low density on 3.5 cm plates before laser treatment. For FISH analysis, a primary fibroblast cell culture derived from a 57, XX sheep goat hybrid was used [6], making it possible to simultaneously map the FISH probes in both species.

2.2. Laser treatment of sheep metaphase chromosomes and PCR amplification

Sheep metaphase chromosomes were spread on a plastic membrane overlaying a 2 cm plate. The metaphases were visualised under a light microscope and a high power laser beam was moved along the chromosome with the exception of the terminal regions according to Metezeau and coworkers [13]. After destruction of the majority of the chromosomes, the remaining DNA material from 10 metaphases was recovered on the plastic film and amplified by DOP-PCR using the 6-MW primer, CCGACTCGAGNNNNNATGTG [17], and called DATS for DOP-Amplified Telomeric Sequences. PCR was carried out in a MJ thermocycler for a total of 30 cycles. After 5 min at 95 °C, 5 low temperature cycles were performed for 1 min at 94 °C, 1.5 min at 30 °C and 3 min extension at 72 °C with a ramping of 3 min between 30 °C and 72 °C. The samples were then subjected to 25 further cycles of 1 min at 94 °C, 1 min at 62 °C and 3 min at 72 °C, this last time being increased by 1 s per cycle. The resulting material was then labelled by random priming using a Boehringer labelling kit and α -³²PdATP.

2.3. Hybridisation to sheep BACs

Five thousand sheep BAC clones plated on two LB-agar 22 cm × 22 cm trays with 12.5 $\mu\text{g} \cdot \text{mL}^{-1}$ chloramphenicol constituted the basic biological material. The BACs were obtained as previously described [18]. Duplicate Hybond N+ membranes (Amersham) were hybridised with the telomeric probe. Positive clones were ascertained after gridding by a second hybridisation step.

2.4. FISH analysis

After BrdU incorporation during the late S phase, air-dried chromosomal preparations were obtained using standard procedures [5]. Protocols used for FISH analysis were previously described [1]. Chromosomal DNA was counterstained and R-banded by the direct fluorescent technique described by Lemieux *et al.* [12]. Competition was performed using 10 μL of sonicated goat DNA (200 ng $\cdot \mu\text{L}^{-1}$) and the prehybridisation was carried out for 2 h 30 min. The slides were screened with a Leica fluorescence microscope and photomicrographs were taken with a Fujichrome 400 Asa colour slide film. Chromosomes were identified using the Texas nomenclature [14].

2.5. Microsatellite isolation and genotyping

BAC DNA was prepared by miniprep as previously described [16]. After complete digestion by *Sau3A1*, BAC fragments were ligated with T4 DNA ligase (Biolabs) in 20 μL of 1X ligase buffer, into 50 ng of the pGEM4Z vector (Promega), linearised by *BamHI* and dephosphorylated using CIAP (Boehringer). Ligations were then diluted to 100 μL , and 1 μL was electroporated in 20 μL DH10B laboratory prepared electrocompetent cells. Each transformation was plated on a 10 cm LB agar dish in the presence of 200 $\mu\text{g} \cdot \text{mL}^{-1}$ ampicillin. After overnight growth, the clones were screened with a mixture of (TG)12 / (TC)12 oligonucleotides fluorescently labelled with the 3'-DIG labelling kit (Boehringer Mannheim). Positive clones were checked by screening after gridding, miniprepped and sequenced using a 377 ABI automatic sequencer. Primers were designed in order to obtain an amplification product in the 100–200 bp range. Genotyping was carried out on ovine reference families [4] by PCR in the presence of α - ^{33}P dATP, using Taq DNA polymerase from Promega or BRL, as previously described [19]. After a denaturation of 5 min at 95 °C, PCR was carried out for 30 cycles (94 °C 15 s, 58 °C 15 s and 72 °C 20 s) in an MJ thermocycler. Samples were loaded on denaturing 5% acrylamide gels, run at 1500 volts for 2 to 3 h. The gels were then dried and autoradiographed.

2.6. Construction of a telomere-enriched library

Total blood was collected on EDTA from sheep by puncture in the jugular vein, and lymphocytes were prepared after lysis of the red cells by NE treatment as previously described [16]. For each plug, 50 μL of NE containing 10^7 cells were mixed with 50 μL of 1% low melting agarose and poured in a 100 μL mold. After solidification, the agarose plug was incubated twice in the lysis buffer (100 mM EDTA, 1% N-lauryl-sarkosyl, and 1 $\text{mg} \cdot \text{mL}^{-1}$ Proteinase K) at 55 °C for 12 h and 4 h. Plugs were then rinsed twice in TE 10 mM–1 mM 30 min). The plugs were then equilibrated 20 min on ice in 2 mL 1X *Bal31* buffer (Promega). The plugs were transferred to 2 mL 1X *Bal31* buffer with 10 μL (30 u) of *Bal31*. Incubation was carried out at 30 °C for 30 min, 120 min and 400 min. In this last case, an additional 10 μL of enzyme was added after 120 min. Plugs were then melted at 65 °C for 15 min, and treated with β -agarase (Boehringer) for 1 h at 45 °C. One phenol/chloroform extraction was then performed by very gentle mixing of the two phases. The upper phase was collected with a cut tip and drop-dialysed for 1 h against 1X *EcoR1* buffer (50 mM Tris-HCl pH 7.5, 100 mM NaCl and 10 mM MgCl_2) using 0.025 μ dialysing membranes (Millipore). After dialysis, the drop was collected in an Eppendorf tube and 3 μL *EcoR1* (30 u) was added for overnight digestion at 37 °C. The DNA was then phenol/chloroform extracted, precipitated and resuspended in TE 10–1 at 160 $\text{ng} \cdot \mu\text{L}^{-1}$, based upon agarose gel estimation. For vector preparation, pGEM4Z was first totally digested with *Sma1*, phenol/chloroform extracted, precipitated and totally digested by *EcoR1*. The double-digested vector was loaded on a preparative 1% agarose gel, and the vector was shortly visualised under 320 nm UV light, cut out of the gel purified by the phenol/liquid nitrogen technique, precipitated and resuspended in TE

10–1 at $50 \text{ ng} \cdot \mu\text{L}^{-1}$. Ligation was performed in $20 \mu\text{L}$ 1X Biolabs ligase buffer, between $1 \mu\text{L}$ *Sma*I/*Eco*R1 vector (50 ng), $1 \mu\text{L}$ of telomere-enriched insert (160 ng), with 80 u of T4 DNA ligase (Biolabs, $0.2 \mu\text{L}$), for 36 h at 8°C . Ligation products were diluted 5 times and $1 \mu\text{L}$ was electroporated in $20 \mu\text{L}$ home made electrocompetent DH10B cells. Transformed bacteria were plated on LB-agar plates with $200 \mu\text{g} \cdot \text{mL}^{-1}$ ampicillin, and screened with a $(\text{CCCTAA})_7$ oligonucleotide γ - ^{32}P -end-labelled with T4 polynucleotide kinase (Eurogentec). Positive clones were gridded and re-screened together with a positive and negative control $\{(\text{T TAGGG})_7$ and $(\text{CCCTAA})_7$ oligonucleotides, respectively}. After this second screening step, positive clones were partially sequenced from both sides of the insert with an ABI377 sequencer.

2.7. BAC library screening and FIGE

Screening was performed by PCR as previously described [16,18]. BAC DNA was prepared by miniprep [16] for further use in FISH or subcloning experiments. Size evaluation of BAC inserts was carried out by Field Inverted Gel Electrophoresis on a FIGE mapper apparatus (BioRad). Briefly, 400 ng (about 1/5 of a miniprep) of BAC DNA was digested by *Not*I in $20 \mu\text{L}$ for 2–3 h and loaded on a 1% agarose gel together with the MidRange and LowRange DNA molecular size marker (BioLabs). The run was performed in the presence of Ethidium Bromide at 14°C for 18 h at 110 V N/S, 170 V S/N with a linear time gradient from 3 s to 15 s.

3. RESULTS

3.1. BAC screening with DATS

BAC plates were leftovers from the construction of the sheep BAC library [18]. The BACs in these plates were not included in the main library because their individual sizes (evaluated by Field Inverted Gel Electrophoresis) were not consistent with the expected size from the size selection experiment, suggesting that they might have been subjected to internal rearrangements. Three $22 \text{ cm} \times 22 \text{ cm}$ trays (about 1700 white colonies per tray) were thus exploited by screening with subtelomeric sequences. Sixty BACs amongst 5000 yielded a positive signal after hybridisation with the DATS and were used in FISH experiments on a sheep/goat hybrid cell line [6]. All the BACs hybridised to the chromosomes of both species at homologous locations. Twenty-three resulted in at least one telomeric or subtelomeric signal (Tabs. I and III, Figs. 1 and 2). Amongst these, 10 BACs hybridised to only one chromosome location, four to three chromosome locations while the 9 other resulted in two signals. Not all the chromosome ends were recognised by the 23 BACs, but several BAC yielded hybridisation signals at the same location. The labelled chromosome ends were 1p-1q, 2q, 3q, 7, 8, 9, 17, 11, 16, 20, 22, 26 using the sheep nomenclature, indicating that 13 extremities amongst 30 were labelled by the isolated BACs. The 14q14 (corresponding to goat or bovine 18q14) internal (non-telomeric) chromosomal region was labelled by two BACs (BAC13 and

Table I. FISH mapping results/library screening with PCR markers.

BAC number	Marker name (EPCDV are microsatellites)	Address in the ordered library	Localisation with BACs from the ordered library	Consistency Genetic vs. Cytogenetic
46	<i>EPCDV001</i>	251e12	1q46(telomeric)	OK
8	<i>EPCDV002</i>	91c3	17q26(telomeric)	OK
8	<i>EPCDV003</i>	74e10	24q14	OK
8	<i>EPCDV004</i>	74e10	24q14	OK
14	<i>EPCDV005</i>	461c6		nd
16	<i>EPCDV006</i>	multiple bands		nd
17	<i>EPCDV007</i>	346d5	2p12	
2	<i>EPCDV008</i>	4g8	9q12	OK
8	<i>EPCDV009</i>	91c3	17q26(telomeric)	OK
15	<i>EPCDV010</i>	37g7 and 46g7	1p43(subtelomeric)	OK
15	<i>EPCDV011</i>	29b5 and 31e6	14q13/14q13-14	OK
13	<i>EPCDV012</i>	31e6	14q13-14	OK
18	<i>EPCDV013</i>	101f6	1q46(telomeric)	OK
19	<i>EPCDV015</i>	252g7	24q21(telomeric)	OK
26	<i>EPCDV016</i>	436c8	8q27(telomeric)	inconsistent
14	<i>EPCDV017</i>	160a6	12q21	
15	<i>EPCDV018</i>	149f5 and 164f5	15q12/3q34	OK
15	<i>EPCDV019</i>	447c8	14q14	OK
24	<i>EPCDV020</i>	272e7	22q25(telomeric)	OK
13	<i>EPCDV021</i>	158d5	1p37(subtelomeric)	OK
15	<i>EPCDV022</i>	158d5	1p37(subtelomeric)	OK
21	<i>EPCDV023</i>	33d2	11q32(subtelomeric)	OK
be1	BE1	31g4	all centromeres	irrelevant
be2	BE1	237b9	all centromeres	irrelevant
be7	BE1	151g6	9q34(telomeric)	irrelevant
be8	BE1	165g7	12q16	irrelevant
be9	BE1	189f9	1p37(telomeric)	irrelevant
be10	BE1	575b6	No signal	irrelevant
be3	BE1	590h11	10q12	irrelevant
be4	BE1	596h12	1p12	irrelevant
be11	BE1	603e6	1p31	irrelevant
be5	BE1	629a11	3q23-25	irrelevant
be6	BE1	648a11	xq38(telomeric)	irrelevant

BAC15). Several other internal chromosome regions were recognised by one BAC (Tab. I).

3.2. Genetic analysis

Twenty-three microsatellites called *EPCDV001* to *EPCDV023* were isolated from 17 out of the 23 telomeric BACs (Tabs. I and II, Fig. 2). No microsatellite could be identified for six BACs. Primers were designed for each marker and

Table II. Characteristics of the microsatellites of the study (telomeric in gray boxes).

(continued on the next page)

Microsatellite	Primer names	Primer sequences	Repeated unit	Size of the amplification product	No. of alleles in the IMF
<i>EPCDV001</i>	EPCDV001.1	GTTGCCTCTCAAAGTGCCTC	(TG)18	180 bp	5
	EPCDV001.2	GGCCTTATGGAAGATTTATGCT			
<i>EPCDV002</i>	EPCDV002.1	CTAGAACTAAGACCAAGCAC	(AC)11	140 bp	2
	EPCDV002.2	CTTTATGATTTGCCCCATATC			
<i>EPCDV003</i>	EPCDV003.1	CAAGAACACTGCTGGAATCAC	(TG)20	155 bp	7
	EPCDV003.2	GAGAGGTCTTGTGACCTATGA			
<i>EPCDV004</i>	EPCDV004.1	CAAGGGGTAACAAAGAGTCA	(CA)10	120 bp	4.1 ⇒ 2 4.2 ⇒ 3
	EPCDV004.2	GAAAGGTAGGCAGCAAGAG			
<i>EPCDV005</i>	EPCDV005.1	GACTTAGCCTATAGCATGAGA	(CA)3(TA)(CA)6	165 bp	Monomorphic
	EPCDV005.2	CAAAGCCTACGTGTATTCAGAG			
<i>EPCDV006</i>	EPCDV006.1	GTCCTCCTGGTTTGCAAAC	(CA)14	185 bp	4
	EPCDV006.2	CTACTGTCTCTTGCTGGGT			
<i>EPCDV007</i>	EPCDV007.3	GATCTGAAACGTGAAGGGTG	(TG)9	130 bp	2
	EPCDV007.2	GCACTCTAGTATTCTTGCCA			
<i>EPCDV008</i>	EPCDV008.1	GACTTCCAAAGAGCTAAGCG	(GT)12	130 bp	9
	EPCDV008.2	GATCTCCTCTAAGCTCACAC			
<i>EPCDV009</i>	EPCDV009.1	GAGCTTTAAGGTTCCAOGA	(GT)6(GA)(GT)10	145 bp	5
	EPCDV009.2	CTTAACCTTCTGGAGGTGAC			
<i>EPCDV010</i>	EPCDV010.3	GCGCACTGGTCTGTCTCC	(CT)16(CA)6	140 bp	6
	EPCDV010.2	CTGACTTGGTTCGAACTTGAG			
<i>EPCDV011</i>	EPCDV011.1	GCATTGAGAAAAGAGACTTGC	(TG)17	160 bp	9
	EPCDV011.2	CACTATGGGAAACAGTATGGA			
<i>EPCDV012</i>	EPCDV012.1	GCAGAGATAATCAGAGCTGC	(TG)24	140 bp	11
	EPCDV012.2	CTAAGTAAGACCTGGCTCCT			

Telomeres in sheep

Table II. *continued.*

Microsatellite	Primer names	Primer sequences	Repeated unit	Size of the amplification product	No. of alleles in the IMF
<i>EPCDV013</i>	EPCDV013.1	CCATCTGTGCTGTCAGTCA	(TG)15	140 bp	5
	EPCDV013.2	GAGCCTGGTGGGTTATAGT			
<i>EPCDV015</i>	EPCDV015.3	CGTGATTGTCCTGCTTGA	((CA) ₄₋₁₄) ₄	120 bp	15.1 → 4 15.2 → 6
	EPCDV015.4	AGGAAAATCACGTCAATGTGT			
<i>EPCDV016</i>	EPCDV016.1	CTCCCCTTCATGCATTCTTG	(TG)16	190 bp	No amplification
	EPCDV016.2	CAGTGTGGTATCTAATCCAGC			
<i>EPCDV017</i>	EPCDV017.1	GCACCCCAAGGTTTATTACA	(TG)17	200 bp	6
	EPCDV017.2	CATCCTTGTTGCTGCATATTG			
<i>EPCDV018</i>	EPCDV018.1	GATTCAGCCACTATGGGAAAC	(CA)15	210 bp	No amplification
	EPCDV018.2	CAAGCTTGCATGCCTGCAGGT			
<i>EPCDV019</i>	EPCDV019.1	CCAGAAATGCCTTCCTTCTG	(TG)14	140 bp	8
	EPCDV019.2	GTCTCTGAGGAGTCACATCT			
<i>EPCDV020</i>	EPCDV020.1	GTGTTTACCATTAGAAATTCC	(AC)16	185 bp	3
	EPCDV020.2	GCCACAGATGACAGGATTG			
<i>EPCDV021</i>	EPCDV021.1	GATCTGCGCACTGCTCTGTC	(CT)15	135 bp	7
	EPCDV021.2	GGCTGACCTGGTCCAACCTG			
<i>EPCDV022</i>	EPCDV022.1	CTGGGATGAGAGACTTCTTAG	(GT)17	165 pb	7
	EPCDV022.2	GCTCACGACTGAATGACTGA			
<i>EPCDV023</i>	EPCDV023.1	CATCTCCTCCCTCCTAACA	(TG)22	115 bp	7
	EPCDV023.2	CAAGAATGGTTTCTAGGGGG			



Figure 1. Fluorescent *in situ* hybridisation of BAC 17, labelling ovine chromosome 8q28 and the goat homeologous chromosome 9q28.




Figure 2. A schematic view of the genetic localisation of microsatellite markers on ovine chromosomes. Double lines indicated markers that have been mapped consistently with cytogenetical data either from the first localisation or after the BAC isolated from the ordered library with the microsatellite primers (see text).

called *EPCDVXXX.1* and *EPCDVXXX.2*. The markers were then genotyped on the international panel of sheep reference families. Genotypes were analysed, making it possible to position them relative to other markers of the sheep International Mapping Flock (IMF) daTab.ase (Tab. III). All were mapped except *EPCDV005* which appeared homozygous on all the parental DNAs, and *EPCDV014*, whose radioactive amplification pattern was too difficult to score (Fig. 2, Tab. IV). Two markers, *EPCDV004* and *EPCDV015* yielded two polymorphic PCR amplification products of different sizes that could be interpreted in a Mendelian transmission. For *EPCDV015*, the two microsatellites were positioned at the end of OAR24, suggesting the existence of a duplication event at this chromosome end. *EPCDV004A* and *4B*, as *EPCDV015A* and *15B* resulted in two signals of different sizes that could be genotyped independently. The distance of 3 centimorgans between *15A* and *15B* (Tab. IV) suggests that duplication probably occurred at the end of ovine chromosome 24. *EPCDV003* and *EPCDV004* originated from the same BAC, the co-localisation of *EPCDV003* and *EPCDV004B* to OAR24 indicates that only the *4B* genotypes are specific of the relevant locus. The other signal *4.A* that was bi-allelic presumably corresponds to an RAPD-type marker. In several cases, the cytogenetical and genetic locations were not consistent, suggesting a high level of chimerism. For instance, BACs 16 and 17 mapped to the telomeric region of sheep chromosome 2q (homeologous to goat chromosomes 2q, Tab. III); however, microsatellites *EPCDV006* and *EPCDV007* were genetically mapped to OAR4 and OAR5 (see Fig. 2). Microsatellites were used for screening in

Table III. Cytogenetic localisations of BACs from the two experiments (telomeric in light gray boxes, non telomeric in black boxes).

Probe number	Size in Kb	Microsatellite(s)	FISH mapping in sheep	FISH mapping in goats
BAC Telo 63	45	-	1q45-46	1q45-46
BAC Telo 46	35	1	2q45	2q45
BAC Telo 01	105	-	3q16-37	3q16-37
BAC Telo 18	130	13	5q21	5q21
BAC Telo 19	110	14,15	5q24-35	5q24-35
BAC Telo 16	40	6	6q22	6q22
BAC Telo 17	130	7	7q13	7q13
BAC Telo 04	70	-	7q24-25	7q24-25
BAC Telo 02	110	8	9q25	9q25
BAC Telo 08	110	2,3,4,9	9q27-28	9q27-28
BAC Telo 14	80	5,17	10q34-35	10q34-35
BAC Telo 15	30	10,11,18,19,22	5	5
BAC Telo 13	80	12,21	12q15	12q15
BAC Telo 28	25	-	14q17-19	14q17-19
BAC Telo 37	65	-	9	9
BAC Telo 26	30	16	14q26	14q26
BAC Telo 24	90	20	16q21	16q21
BAC Telo 45	25	-	17q23-2	17q23-2
BAC Telo 59	25	-	4	4
BAC Telo 21	80	23	18q14	18q14
BAC Telo 25	25	-	19q22	19q22
BAC Telo 09	65	-	20q22	20q22
BAC Telo 05	80	-	23q25	23q25
			26q23	26q23
			27q24	27q24
			14q34	14q34
			3q37	3q37
			Xq38	Xq38

Telomeres in sheep

order to assess the chimerism of the BACs used in the screening experiment, compared to the bulk of the library which has been stored and ordered [18]. BAC addresses are reported in Table I. Contrarily to the first series of BACs obtained after the screening with DATs, all these BACs from the ordered library resulted in a unique signal in the FISH experiments (Tab. I).

Table IV. Two-points linkage tests.

Marker	Locus	Min Rec	Max Rec	Max Lod	θ	Lod
<i>EPCDV001</i>	<i>HH36</i>	8	123	26.355	0.06	22.769
	<i>BMS2263</i>	3	100	25.115	0.03	22.405
	<i>URB14H</i>	17	121	19.173	0.12	15.915
	<i>McM357</i>	9	98	18.794	0.08	15.815
	<i>UWCA46</i>	26	96	9.277	0.22	6.971
	<i>BMS599</i>	14	71	9.072	0.17	7.047
	<i>BM1824</i>	34	93	6.187	0.28	3.534
	<i>MAF4</i>	38	94	5.326	0.30	3.034
<i>EPCDV002</i>	<i>UCDO48</i>	0	8	2.408	0.00	2.107
<i>EPCDV003</i>	<i>ILSTS43</i>	6	119	27.174	0.05	24.161
	<i>BMS2843</i>	13	132	24.648	0.09	21.939
	<i>BP28</i>	19	147	24.325	0.11	21.007
	<i>BM4005</i>	11	120	23.029	0.08	20.317
	<i>BM737</i>	23	149	22.391	0.13	18.774
	<i>BMS744</i>	19	120	17.762	0.14	14.450
	<i>INR206</i>	11	93	16.060	0.11	13.648
	<i>B1864A</i>	3	53	11.777	0.05	10.269
	<i>ILSTS102</i>	34	122	11.439	0.22	8.153
	<i>JMP29</i>	32	105	8.900	0.23	5.908
	<i>SPN</i>	12	63	8.256	0.16	7.052
	<i>PRM2</i>	3	39	7.950	0.07	7.648
	<i>ALDOA</i>	17	70	7.526	0.20	5.431
	<i>EP0042</i>	0	22	6.623	0.00	6.021
	<i>BMS514</i>	20	70	6.388	0.23	3.535
	<i>UCDO12</i>	7	42	6.023	0.14	5.119
<i>EPCDV004.A</i>	<i>INR108</i>	0	34	10.235	0.00	9.633
	<i>BM746</i>	1	36	9.142	0.03	8.236
	<i>TGLA77</i>	1	36	9.142	0.03	8.236
	<i>FCB129</i>	1	35	8.853	0.03	7.961
	<i>TGLA58</i>	4	33	5.634	0.11	4.781
	<i>BMS710</i>	4	33	5.634	0.11	4.781
	<i>BM8118</i>	1	22	5.137	0.04	4.534
	<i>BMS1915</i>	3	28	5.052	0.10	4.148
	<i>BMS460</i>	3	27	4.795	0.10	3.898
	<i>CP34</i>	6	31	4.016	0.16	3.115
	<i>TGLA67</i>	2	20	3.712	0.09	3.411

Marker	Locus	Min Rec	Max Rec	Max Lod	θ	Lod
<i>EPCDV004.2</i>	<i>EPCDV009</i>	0	22	6.623	0.00	6.021
	<i>BMS2843</i>	2	20	3.712	0.09	3.110
	<i>ILSTS43</i>	1	16	3.466	0.06	2.864
	<i>INR209</i>	0	11	3.311	0.00	3.010
	<i>INR206</i>	0	11	3.311	0.00	3.010
	<i>INR129</i>	0	11	3.311	0.00	3.010
<i>EPCDV006</i>	<i>MAF70</i>	3	116	29.741	0.03	27.010
	<i>McM144</i>	8	85	16.152	0.09	14.082
	<i>CRSP13</i>	7	64	11.445	0.10	9.639
	<i>RM67</i>	4	54	11.138	0.07	9.633
	<i>BM6437</i>	7	54	8.923	0.12	6.815
	<i>MAF50</i>	4	42	7.945	0.09	6.740
	<i>IL6 PM</i>	12	58	7.144	0.17	5.063
	<i>BMS1237</i>	4	38	6.907	0.10	6.001
<i>EPCDV007</i>	<i>TGLA176</i>	0	58	17.460	0.00	15.955
	<i>McM380</i>	2	56	13.682	0.03	12.168
	<i>RD2138</i>	3	48	10.397	0.06	8.892
	<i>URB051</i>	0	26	7.827	0.00	7.225
	<i>IL5</i>	1	24	5.702	0.04	5.100
	<i>BM9289</i>	0	18	5.419	0.00	4.816
	<i>TG303</i>	3	29	5.309	0.09	4.405
	<i>RM6</i>	1	20	4.576	0.05	3.974
	<i>IL3</i>	4	26	3.915	0.13	3.011
<i>EPCDV008</i>	<i>MAF33</i>	15	146	26.804	0.09	23.529
	<i>BM4630</i>	13	124	22.576	0.10	19.302
	<i>BL1009</i>	24	142	20.183	0.14	16.868
	<i>CSPS11</i>	16	113	17.831	0.12	15.419
	<i>HH29</i>	8	88	16.940	0.08	14.836
	<i>ILSTS11</i>	17	99	13.928	0.15	11.589
	<i>MS1678</i>	16	91	12.605	0.15	10.801
	<i>CSSM55</i>	16	80	10.114	0.17	7.423
	<i>ILSTS8</i>	11	67	9.699	0.14	7.303
	<i>RD240</i>	40	113	7.880	0.26	5.214
	<i>CSSM66</i>	41	112	7.436	0.27	4.829
	<i>McMA10</i>	21	76	7.191	0.22	4.796
	<i>BMS740</i>	37	95	5.727	0.29	2.992
<i>EPCDV009</i>	<i>VNTR11</i>	11	113	21.197	0.09	18.491
	<i>ILSTS58</i>	23	120	15.656	0.16	12.721
	<i>McMA20</i>	19	107	14.723	0.15	12.085
	<i>BM8125</i>	22	98	11.295	0.18	8.972
	<i>BM1862</i>	6	57	10.360	0.10	7.995
	<i>BM7136</i>	34	112	9.538	0.24	6.696
	<i>HUJ223</i>	16	71	8.157	0.18	6.951

Marker	Locus	Min Rec	Max Rec	Max Lod	θ	Lod
	<i>TG322</i>	21	80	7.981	0.21	5.977
	<i>URB002</i>	28	91	7.626	0.24	5.382
	<i>EPCDV002</i>	0	25	7.526	0.00	6.924
<i>EPCDV010</i>	<i>BMS2833</i>	9	107	21.175	0.08	19.065
	<i>INR197</i>	12	109	19.437	0.10	16.728
	<i>BMS835</i>	23	116	14.761	0.17	11.754
	<i>TGLA127</i>	5	54	10.325	0.09	8.596
	<i>INRA88</i>	5	50	9.280	0.09	7.173
	<i>RM309A</i>	24	92	9.236	0.21	6.629
	<i>HU1177</i>	36	103	7.313	0.26	4.638
	<i>BMS2145</i>	12	58	7.144	0.17	5.062
	<i>HH51</i>	39	103	6.495	0.28	3.582
	<i>BM4301</i>	36	98	6.473	0.27	3.816
	<i>McMA41</i>	37	95	5.727	0.28	3.076
	<i>BM3020</i>	34	87	5.216	0.29	2.906
<i>EPCDV011</i>	<i>BMS2213</i>	4	152	38.882	0.03	35.547
	<i>MT2</i>	18	140	23.228	0.11	19.911
	<i>S1915b</i>	16	132	22.535	0.11	19.524
	<i>RD270</i>	18	133	21.497	0.12	18.487
	<i>UWCA28</i>	14	94	14.421	0.13	11.112
	<i>INR138</i>	28	119	13.166	0.19	9.866
	<i>HAUT14</i>	36	125	11.308	0.23	8.042
	<i>MS2355</i>	18	82	9.631	0.18	6.927
	<i>TEX10</i>	6	53	9.336	0.10	7.576
	<i>McMA17</i>	21	84	8.789	0.20	6.390
	<i>TGLA357</i>	35	109	8.666	0.24	5.959
	<i>McM133</i>	21	83	8.586	0.20	6.181
	<i>BM7109</i>	40	107	6.882	0.28	4.041
	<i>INRA63</i>	48	119	6.768	0.30	3.555
	<i>CSRD232</i>	44	105	5.586	0.30	3.057
<i>EPCDV012</i>	<i>BMS2213</i>	2	162	44.678	0.01	41.350
	<i>EPCDV011</i>	6	151	36.200	0.04	32.886
	<i>MT2</i>	20	147	23.695	0.12	20.383
	<i>RD270</i>	14	129	23.146	0.10	20.435
	<i>S1915b</i>	14	126	22.379	0.10	19.669
	<i>UWCA28</i>	10	96	17.525	0.09	14.209
	<i>INR138</i>	23	116	14.761	0.17	11.754
	<i>McMA17</i>	17	100	14.160	0.15	11.450
	<i>TEX10</i>	5	67	13.788	0.07	11.713
	<i>HAUT14</i>	39	138	12.746	0.22	9.174
	<i>TGLA357</i>	32	120	11.783	0.21	9.076
	<i>McM133</i>	23	87	8.618	0.21	6.215
	<i>BMS2355</i>	17	73	8.151	0.19	5.751

Marker	Locus	Min Rec	Max Rec	Max Lod	θ	Lod
	<i>BM7109</i>	46	109	5.725	0.30	3.243
<i>EPCDV013</i>	<i>URB14H</i>	5	148	36.493	0.03	33.475
	<i>McM357</i>	3	94	23.388	0.03	20.980
	<i>HH36</i>	7	102	21.526	0.06	18.809
	<i>BMS2263</i>	11	97	17.074	0.10	14.966
	<i>EPCDV001</i>	19	109	15.185	0.15	12.475
<i>EPCDV015.A</i>	<i>EPCDV015.2</i>	3	105	26.558	0.03	24.446
	<i>RM24L</i>	4	49	9.796	0.08	9.190
	<i>ELN</i>	13	63	7.776	0.17	6.573
	<i>EL01</i>	2	33	7.207	0.06	6.302
	<i>ALDOA</i>	9	47	6.136	0.16	4.947
	<i>BM737</i>	33	77	3.931	0.31	2.038
<i>EPCDV015.B</i>	<i>EPCDV015.1</i>	3	105	26.558	0.03	24.446
	<i>RM24L</i>	3	59	13.447	0.05	12.543
	<i>ELN</i>	14	62	7.111	0.19	5.699
	<i>EL01</i>	6	38	5.634	0.14	4.431
	<i>BM737</i>	31	81	5.022	0.28	2.787
	<i>ALDOA</i>	11	46	5.016	0.19	4.112
	<i>HH72</i>	6	32	4.241	0.16	3.940
	<i>BP28</i>	37	80	3.513	0.32	1.804
	<i>UCDO12</i>	0	11	3.311	0.00	3.010
<i>EPCDV016</i>	<i>BMS1967</i>	3	80	19.381	0.04	17.868
	<i>URB024</i>	6	70	13.762	0.08	12.257
	<i>BM3215</i>	7	63	11.189	0.10	9.985
	<i>McMA42</i>	1	38	9.720	0.03	8.511
	<i>McM494</i>	17	72	7.942	0.19	6.159
	<i>BM4208</i>	19	68	6.358	0.22	4.886
	<i>RD275</i>	3	29	5.309	0.10	3.852
<i>EPCDV017</i>	<i>BMS2258</i>	7	123	27.295	0.05	24.276
	<i>BMS792</i>	3	93	23.101	0.03	20.993
	<i>CRSP14</i>	20	86	9.614	0.19	7.512
	<i>TG137</i>	23	91	9.423	0.20	7.115
	<i>INR192</i>	0	26	7.827	0.00	7.225
	<i>McM527</i>	34	99	7.203	0.26	4.519
	<i>BM1853</i>	3	36	7.147	0.08	5.677
	<i>UCDO37</i>	0	11	3.311	0.00	3.010
<i>EPCDV018</i>	<i>BR3510</i>	8	52	7.830	0.13	6.624
	<i>BMS1004</i>	14	45	3.721	0.24	2.259
	<i>BMS1MDR</i>	0	12	3.612	0.00	3.311
	<i>McMA16</i>	0	11	3.311	0.00	3.010
	<i>UCDO37</i>	0	11	3.311	0.00	3.010
	<i>BM4025</i>	16	45	3.118	0.27	1.968

Marker	Locus	Min Rec	Max Rec	Max Lod	θ	Lod
	<i>McMA49</i>	0	10	3.010	0.00	2.709
<i>EPCDV019</i>	<i>EPCDV012</i>	3	115	29.451	0.03	27.023
	<i>BMS2213</i>	1	102	28.561	0.01	26.454
	<i>EPCDV011</i>	4	96	22.809	0.04	20.702
	<i>MT2</i>	13	105	17.746	0.11	15.337
	<i>RD270</i>	10	84	14.462	0.11	12.653
	<i>S1915b</i>	12	87	13.922	0.12	11.815
	<i>McMA17</i>	11	72	10.885	0.13	9.080
	<i>HAUT14</i>	22	96	10.871	0.19	8.463
	<i>UWCA28</i>	4	50	10.063	0.07	7.953
	<i>TEX10</i>	3	44	9.303	0.07	7.823
	<i>TEX10</i>	3	42	8.760	0.07	7.284
	<i>TG357</i>	22	81	7.804	0.21	5.999
	<i>INR138</i>	18	70	7.128	0.21	5.327
	<i>EPCDV020</i>	<i>BMS2721</i>	4	71	15.795	0.05
<i>TG272</i>		4	53	10.869	0.07	9.364
<i>McM156</i>		7	56	9.421	0.11	8.216
<i>BMS2614</i>		12	63	8.256	0.16	6.751
<i>BMS1620</i>		9	56	8.214	0.14	6.720
<i>AE64</i>		8	50	7.354	0.14	6.150
<i>MS2466</i>		5	41	6.980	0.11	6.077
<i>ILSTS20</i>		7	42	6.023	0.14	4.819
<i>McM139</i>		7	41	5.790	0.15	4.299
<i>ILSTS5</i>		6	38	5.634	0.14	5.031
<i>BMS823</i>		5	31	4.537	0.14	3.935
<i>FNTBEB</i>		2	23	4.499	0.08	2.998
<i>EPCDV021</i>		<i>EPCDV010</i>	7	127	28.405	0.05
	<i>INR197</i>	14	104	16.857	0.12	14.148
	<i>BMS2833</i>	12	96	16.150	0.11	14.344
	<i>BMS835</i>	21	108	13.943	0.16	11.233
	<i>PPT</i>	3	54	12.054	0.05	10.849
	<i>RM309A</i>	21	86	9.200	0.20	6.887
	<i>HU1177</i>	32	97	7.448	0.25	5.048
	<i>TG127</i>	6	45	7.330	0.12	5.935
	<i>BMS2145</i>	11	53	6.512	0.17	4.728
	<i>HH51</i>	36	96	6.145	0.28	3.507
	<i>EPCDV022</i>	<i>EPCDV010</i>	4	125	31.089	0.03
<i>EPCDV021</i>		9	110	21.974	0.08	19.258
<i>BMS2833</i>		11	105	19.123	0.09	16.707
<i>INRA197</i>		11	95	16.566	0.10	13.854
<i>BMS835</i>		22	109	13.685	0.17	10.392
<i>TG127</i>		5	52	9.801	0.09	7.816
<i>PPT</i>		4	48	9.529	0.08	8.324

Marker	Locus	Min Rec	Max Rec	Max Lod	θ	Lod
	<i>INRA88</i>	6	50	8.577	0.11	6.175
	<i>RM309A</i>	23	81	7.443	0.22	5.146
	<i>BMS2145</i>	13	54	5.853	0.20	3.529
	<i>HH51</i>	41	97	5.080	0.31	2.033
<i>EPCDV023</i>	<i>McM120</i>	24	146	21.119	0.14	18.186
	<i>CSSM15</i>	22	129	18.229	0.15	14.986
	<i>CSSM65</i>	7	86	17.209	0.08	15.698
	<i>MB087</i>	2	66	16.551	0.03	13.541
	<i>TAU</i>	2	66	16.551	0.03	13.541
	<i>McMA24</i>	7	74	14.035	0.09	12.828
	<i>CRSP6</i>	38	140	13.498	0.22	10.127
	<i>McMA29</i>	19	101	13.355	0.16	10.150
	<i>BM17132</i>	16	77	9.453	0.17	7.063
	<i>BMS501</i>	1	30	7.413	0.03	5.606
	<i>ETH3</i>	6	45	7.330	0.12	5.824
	<i>RD281h</i>	16	67	7.315	0.20	4.757

3.3. Cloning of sheep subtelomeric sequences

The enrichment protocol was based upon a selection by the vector of inserts carrying *Eco*R1 cohesive ends on one side and blunt ends on the other side (Fig. 3). Two thousand clones of telomere enriched DNA were plated, and screened with the (TTAGGG)₇ probe, making it possible to select 18 positive clones (about 1%). The size of the DNA insert was evaluated by agarose gel electrophoresis after *Hind*III digestion and was comprised between 1 and 15 kb (average ~ 4 kb). Partial sequencing was carried out from both sides and revealed telomeric repeats on the reverse sequencing side (*Sma*I) for a clone called BE1. Primers were designed from the *Eco*R1 sequence in order to screen the sheep BAC library. The screening with BE1 primers indicated that 20 superpools were positive amongst the 39 of the library, suggesting that BE1 primers recognise repetitive elements in sheep DNA. Whilst it was not possible to identify BAC addresses from the strongest responses in the superpool, increasing the PCR specificity by adding 2% formamide and raising the temperature to 60 °C made it possible to recover 11 different BACs from the ordered BAC library (be1 to be11, Tab. I). These positive BAC clones were then mapped by FISH (Tabs. I and III). Two resulted in centromeric signals, one did not give any clear response and five were inside chromosome arms. The three others (be6, be7) yielded strong signals at the telomeres of chromosomes X, 9 and 1, respectively.

4. DISCUSSION

In this study, two techniques are presented that allow the identification of subtelomeric sequences in sheep, where such sequences have not been cloned




Figure 3. A scheme of the telomere enrichment procedure. Lymphocytes were included in low melting agarose prior to DNA extraction with Proteinase K. The enrichment is based upon the fact that after treatment, unfragmented chromosomes will result in *EcoRI*/*EcoRI* fragments or in blunt/*EcoRI* fragments, if they encompass a telomeric region. This technique results in a 100 X enrichment in such sequences over an *EcoRI* library (see text).

before. The first technique was based upon the screening of a sheep BAC library with telomeric and subtelomeric material preserved from high-energy laser light and DOP-PCR amplified. Since this technique does not give clear information about the distance between the BACs obtained and the true physical chromosome end, another complementary approach was designed. Chromosome ends

have already been cloned in humans using a modified YAC vector containing a unique telomere, where the second chromosome telomere is provided by the genomic DNA [3]. By contrast, the direct cloning of telomere sequences is impossible in classical BAC vectors because the genomic DNA is prepared by partial digestion of high molecular weight DNA. For instance, in our sheep library, *Hind*III has been used as the cloning enzyme. Consequently, all the BAC inserts end with a *Hind*III restriction site, implicating an impossibility to clone the chromosome end, made of TTAGGG repeats. To circumvent this problem, we decided to identify inside the ordered sheep BAC library [18], the BAC(s) containing the last *Hind*III site of the chromosome. For this, a telomere-enriched library was constructed, with clones large enough to encompass the last *Hind*III site. The particularity of this library is that the telomeric side of the clones is blunt ended and shortened by the *Bal*31 exonuclease, while the other side is terminated by an *Eco*R1 site, generally rarer than *Hind*III sites in mammalian genomes. There is thus a good probability that the last *Hind*III site will be comprised in the telomere-enriched DNA, implying that a side of the insert is present in the BAC library. *Bal*31 is used in such a way that not all the telomeric (TTAGGG) repeats are digested, making it possible to screen the enriched library by hybridisation with specific telomeric sequences. The sequence from the *Eco*R1 side makes it possible to choose primers, to screen the sheep BAC library, and to obtain the BAC terminating the chromosome. About 1% of positive clones could be identified in the TED, making it possible to evaluate the enrichment achieved in this library. Indeed, a medium sheep chromosome contains 100 Mb of DNA, which when digested by *Eco*R1 (1 site every 4 000 bp of DNA) yield on average 25 000 fragments, of which two (about 0.01%) are telomeric. Consequently, the protocol designed in this paper results in a 100 X enrichment, and could be efficiently applied to other organisms, where subtelomeric regions deserve further analysis.

BAC clones were identified by the first technique and 23 actually mapped to subtelomeric regions. The hybridisation of 37 other BACs to internal chromosome regions can be explained either by the presence of telomere-like sequences inside chromosomes, or by the non-complete destruction of the non-telomeric sequence by the laser treatment. Often, the 23 BACs hybridising a telomeric region recognised more than one telomere, and sometimes also some non-telomeric regions. These results have been confirmed by the genetic analysis performed after genotyping the microsatellites isolated in the sheep reference families [4]. The hybridisation to different chromosomal regions suggests several possible explanations: either the BACs used were unusually chimerical (whilst it is assumed that the chimerism rate in BACs is around 5%), or some subtelomeric sequences were homologous between different chromosomes, leading to different hybridisation signals. FIGE analysis of the BACs from the first experiment (Tab. I) made it possible to estimate insert sizes between 25 kb and 130 kb, with an average of 69 kb (± 35 kb). This figure is about half the average insert size of the clones in the library [18], suggesting that the subset used for screening with DATs was qualitatively different from the rest of the library. A further proof was obtained by screening the ordered library with the different microsatellites isolated, when they yielded hybridisation signals on different chromosomes, such as *EPCDV010* and *EPCDV011*, both isolated

from BAC 15, but genetically mapped to chromosomes 1p and 14, respectively. In all these cases, the two microsatellites belonged to different BACs from the ordered library, substantiating the possibility of a high level of chimerism of the leftover BACs used for screening with DATs. The intrachromosomal regions 14q14 labelled independently by BACs 13 and 15 might be interpreted as the remnant of a "fossil" telomeric region. Actually, the region roughly corresponds to a breakpoint between two human chromosome regions, 16q13 and 16q24 and between mouse regions 8.40 and 8.61, suggesting a possible position for an evolutionary breakpoint [15]. To summarise, this first approach to isolate subtelomeric DNA sequences, was impaired by the low quality of the BACs screened that displayed a high level of chimerism. This was illustrated at the cytogenetic level since different signals were visible in FISH, but also at the genetic level since when different microsatellites were isolated from several of these BACs they happened to be linked to markers from different chromosome locations. By contrast, in the ordered sheep BAC library, the BACs contain very few chimeric inserts. This explains why when the microsatellites originating from a BAC from the first series were used to screen the ordered sheep BAC library by PCR, each microsatellite resulted in one address.

The telomere enrichment technique presented in this paper could be an efficient way to provide telomeric sequences from a wide range of species. The primers from probe BE1 made it possible to identify 11 different BACs including two labelled centromeres, suggesting a high content in repetitive sequences (satellite-type), while three clearly labelled telomeric regions. This approach could be validated in the future by isolating new BE probes. The mapping of microsatellites originating from this material will presumably position them at the ultimate extremity of the chromosomes. They could thus constitute the best markers for following chromosome erosion after cloning experiments in ruminants.

REFERENCES

- [1] Bahri-Darwich I., Vaiman D., Olsaker I., Oustry A., Cribiu E.P., Assignment of bovine synteny groups U27 and U8 to R-banded chromosome 12 and 27, respectively, *Hereditas* 120 (1994) 261-5.
- [2] Barendse W., Vaiman D., Kemp S.J., Sugimoto Y., Armitage S.M., Williams J.L., Sun H.S., Eggen A., Agaba M., Aleyasin S.A., Band M., Bishop M.D., Buitkamp J., Byrne K., Collins F., Cooper L., Coppettiers W., Denys B., Drinkwater R.D., Easterday K., Elduque C., Ennis S., Erhardt G., Li L., *et al.*, A medium-density genetic linkage map of the bovine genome, *Mamm. Genome* 8 (1997) 21-8.
- [3] Brown W.R., Molecular cloning of human telomeres in yeast, *Nature* 338 (1989) 774-6.
- [4] Crawford A.M., Dodds K.G., Ede A.J., Pierson C.A., Montgomery G.W., Garmonsway H.G., Beattie A.E., Davies K., Maddox J.F., Kappes S.W., *et al.*, An autosomal genetic linkage map of the sheep genome, *Genetics* 140 (1995) 703-24.
- [5] Cribiu E.P., Matejka M., Darre R., Durand V., Berland H.M., Bouvet A., Identification of chromosomes involved in a robertsonian translocation in cattle, *Genet. Sel. Evol.* 21 (1989) 555-560.

- [6] Cribiu E.P., Matejka M., Denis B., Malher X., Étude chromosomique d'un hybride chèvre × mouton fertile, *Génét. Sél. Evol.* 20 (1988) 379–386.
- [7] Cross S., Lindsey J., Fantès J., McKay S., McGill N., Cooke H., The structure of a subterminal repeated sequence present on many human chromosomes, *Nucleic Acids Res.* 18 (1990) 6649–57.
- [8] Cross S.H., Allshire R.C., McKay S.J., McGill N.I., Cooke H.J., Cloning of human telomeres by complementation in yeast, *Nature* 338 (1989) 771–4.
- [9] Griffith J.D., Comeau L., Rosenfield S., Stansel R.M., Bianchi A., Moss H., de Lange T., Mammalian telomeres end in a large duplex loop, *Cell* 97 (1999) 503–14.
- [10] Gu F., Hindkjaer J., Primed *in situ* labeling (PRINS) detection of telomeric (CCCTAA)_n sequences in chromosomes of domestic animals, *Mamm. Genome* 7 (1996) 231–232.
- [11] Klobutcher L.A., Swanton M.T., Donini P., Prescott D.M., All gene-sized DNA molecules in four species of hypotrichs have the same terminal sequence and an unusual 3' terminus, *Proc. Natl. Acad. Sci. U S A* 78 (1981) 3015–9.
- [12] Lemieux N., Dutrillaux B., Viegas-Pequignot E., A simple method for simultaneous R or G banding by fluorescence *in situ* hybridization of small single copy genes, *Cytogenet. Cell Genet.* 59 (1992) 311–312.
- [13] Metezeau P., Kiefer H., Lamy G., Delamare G., Analytic and preparative laser scanning cytometry, *Pathol. Biol. (Paris)* 41 (1993) 276–80.
- [14] Popescu C.P., Long S., Riggs P., Womack J., Schmutz S., Fries R., Gallagher D.S., Standardization of cattle karyotype nomenclature: Report of the committee for the standardization of the cattle karyotype, *Cytogenet. Cell Genet.* 74 (1996) 259–261.
- [15] Schibler L., Vaiman D., Oustry A., Giraud-Delville C., Cribiu E.P., Comparative gene mapping: A fine-scale survey of chromosome rearrangements between ruminants and humans, *Genome Res.* 8 (1998) 901–15.
- [16] Schibler L., Vaiman D., Oustry A., Guinec N., Dangy-Caye A.L., Billault A., Cribiu E.P., Construction and extensive characterization of a goat bacterial artificial chromosome library with threefold genome coverage, *Mamm. Genome* 9 (1998) 119–24.
- [17] Telenius H., Carter N.P., Bebb C.E., Nordenskjöld M., Ponder B.A.J., Tunnaccliffe A., Degenerate oligonucleotide primer PCR: general amplification of target DNA by a single degenerate primer, *Genomics* 13 (1992) 718–725.
- [18] Vaiman D., Billault A., Tabet-Aoul K., Schibler L., Vilette D., Oustry-Vaiman A., Soravito C., Cribiu E.P., Construction and characterization of a sheep BAC library of three genome equivalents, *Mamm. Genome* 10 (1999) 585–7.
- [19] Vaiman D., Schibler L., Bourgeois F., Oustry A., Amigues Y., Cribiu E.P., A genetic linkage map of the male goat genome, *Genetics* 144 (1996) 279–305.